

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/79255 A1

- (51) International Patent Classification⁷: C07J 9/00, 21/00, 41/00, 31/00
- (21) International Application Number: PCT/US01/12004
- (22) International Filing Date: 12 April 2001 (12.04.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/196,646 12 April 2000 (12.04.2000) US
- (71) Applicant (for all designated States except US):
GENAERA CORPORATION [US/US]; 5110 Campus Drive, Plymouth Meeting, PA 19462 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **KINNEY, William, A.** [US]; 5110 Campus Drive, Plymouth Meeting, PA 19462 (US). **ZHANG, Xuehai** [US]; 5110 Campus Drive, Plymouth Meeting, PA 19462 (US). **MICHALAK, Ronald** [US]; 5110 Campus Drive, Plymouth Meeting, PA 19462 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
- with international search report
 - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: A PROCESS FOR THE PREPARATION OF 7.ALPHA.-HYDROXY 3-AMINOSUBSTITUTED STEROLS USING INTERMEDIATES WITH AN UNPROTECTED 7.ALPHA.-HYDROXY GROUP



(57) Abstract: An efficient method for the synthesis of aminosterol compounds such as squalamine and compound 1436 is described. A method of the invention provides for regioselective oxidation and regioselective sulfonation of a fused ring system. The fused ring base can be, for example, a steroid ring base. The aminosterol compounds are effective as, among others, antibiotics, antiangiogenic agents and NHE3 inhibitors.

WO 01/79255 A1

**Regioselective and Stereoselective Oxidation of
Fused Ring Systems Useful For The Preparation Of Aminosterols
GOVERNMENT FUNDING**

The research described in this patent application was funded in part by Small
5 Business Innovative Research Grant #1 R43 CA 80473-01 from the National Cancer
Institute of the National Institutes of Health.

BACKGROUND OF THE INVENTION

Field of the Invention

10 The invention relates to a novel method of producing fused ring based
compounds or aromatics including aminosterol compounds. A method of the
invention offers regioselective oxidation and regioselective sulfonation of fused ring
systems with few protecting groups. The aminosterol compounds produced by a
method of the invention are useful as, among others, antibiotics, antiangiogenic
15 agents and NHE3 inhibitors.

Description of the Related Art

Several aminosterol compositions have been isolated from the liver of
the dogfish shark, *Squalus acanthias*. One important aminosterol is squalamine
20 (3 β -(N-[3-aminopropyl]-1,4-butanediamine)-7 α ,24R-dihydroxy-5 α -cholestane
24-sulfate), illustrated Fig. 1. The aminosterol squalamine, which includes a
sulfate group at the C-24 position, is the subject of U.S. Patent No. 5,192,756
which also describes the aminosterol's antibiotic properties.

Since the discovery of squalamine, however, several other interesting
25 properties of this compound have been discovered. For example, as described in
U.S. Patent Nos. 5,733,899 and 5,721,226, squalamine may function as an
antiangiogenic agent useful for the treatment of cancers. See U.S. Patent No.
6,147,060. Additional uses of squalamine such as an agent for inhibiting NHE3
and as an agent for inhibiting endothelial cell growth are disclosed in U.S. Patent
30 No. 5,792,635.

Methods for synthesizing squalamine have been described. See WO

94/19366 which relates to U.S. Patent Application. No. 08/023,347. U.S. Patent No. 5,792,635 also discloses squalamine isolation and synthesis techniques.

Stemming from the discovery of squalamine, other aminosterols have been discovered in the dogfish shark liver and have been investigated. One important
5 aminosterol that has been isolated and identified as "compound 1436" or simply "1436" has the structure shown in Fig. 2. This compound has the general molecular formula $C_{37}H_{72}N_4O_5S$ and a calculated molecular weight of 684.53017. Like squalamine, this aminosterol has a sulfate group at the C-24 position.

Compound 1436 previously has been described in U.S. Patent. No.
10 5,795,885. As further described in this patent, compound 1436 has a variety of interesting properties. For example, compound 1436 inhibits human T-lymphocyte proliferation, as well as the proliferation of a wide variety of other cells and tissues. Additional uses of compound 1436 are disclosed in U.S. Patent 6,143,738. U.S. Patent Nos. 5,795,885 and 5,847,172 also describe the structure of compound 1436
15 as well as processes for synthesizing and isolating the compound. For example, compound 1436 can be prepared from a squalamine starting material.

Difficulties have been encountered, however, when attempting to provide a process for synthesizing squalamine or compound 1436 from commercially available starting materials (i.e., not from shark liver isolates). These difficulties
20 include low overall yields of the desired steroid product as well as multiple synthetic steps.

Additional difficulties are encountered in providing a sulfate group at the C-24 position. Particularly, it is difficult to provide the sulfate group at the C-24 position in a highly stereoselective orientation. See, for example, Pechulis, et al.,
25 "Synthesis of 24R - Squalamine, an Anti-Infective Steroidal Polyamine," *J. Org. Chem.*, 1995, Vol. 60, pp. 5121-5126; and Moriarty, et al., "Synthesis of Squalamine. A Steroidal Antibiotic from the Shark," *Tetrahedron Letters*, Vol. 35, No. 44, (1994), pp. 8103-8106.

Because of the importance of squalamine, compound 1436, other
30 aminosterols, 24R and 24S-hydroxylated steroids and vitamin-D₃ metabolites, there has been considerable interest in preparing stereospecific compounds especially at

the C-24 position. As mentioned above, processes for producing squalamine and compound 1436 have been described. However, these processes do not enable large scale production of the desired aminosterol compounds because relatively low yields are realized by these processes.

Processes for stereoselectively producing cerebrosterol, MC 903, and 1α , 24(R)-dihydroxyvitamin D₃ have been developed. Koch, *et al.*, "A Stereoselective Synthesis and a Convenient Synthesis of Optically Pure (24R)- and (24S)-24 hydroxycholesterols," *Bulletin de la Société Chimique de France*, 1983, (No. 7-8), Vol. II, pp. 189-194; Calverley, "Synthesis of MC 903, a Biologically Active Vitamin D Metabolite Analogue," *Tetrahedron*, 1987, Vol. 43, No. 20, pp. 4609-4619; and Okamoto, *et al.* "Asymmetric Isopropylation of Steroidal 24-Aldehydes for the Synthesis of 24(R)-Hydroxycholesterol," *Tetrahedron: Asymmetry*, 1995, Vol. 6, No. 3, pp. 767-778. These processes attempt to reduce 22-ene-24-one and 22-yne-24-one systems in a stereoselective manner. Unfortunately, the processes were not highly stereospecific and often resulted in mixtures of the 24R and 24S which were difficult to separate. Thus these processes were not conducive to large scale synthesis.

Other attempts were also not conducive to large scale synthesis. These processes suffered from being too lengthy or impractical. For example, successful reduction has been achieved of a related 25-ene-24-one system using Noyori's 2,2'-dihydroxy-1,1'-binaphthyl lithium aluminum hydride reagent at -90°C to give 95:5 selectivity for the 24R-alcohol. Ishiguro, *et al.* "Stereoselective Introduction of Hydroxy-Groups into the 24-, 25-, and 26-Positions of the Cholesterol Side Chain," *J. C. S. Chem. Comm.*, 1981, pp. 115-117. However, the 25-ene-24-one intermediate material (producible in four steps) is less readily accessible than the 22-ene-24-one system (producible in one step). Furthermore, the low temperature required for the chiral reduction also detracts from the commercial practicality of this method.

A large scale stereoselective synthesis has been developed to satisfy the requirements for rapid entry in Phase I clinical trials. Zhang, X., *et al.*, *J. Org. Chem.*, 63, 8599-8603 (1998). However, the synthesis suffered two major

drawbacks. First, the synthesis was quite lengthy. Secondly, introduction of a 7 α -hydroxyl group proved problematic.

Thus there exists a need in the art for a method of preparing aminosterol compounds such as squalamine, compound 1436 and various homologs that
5 overcome the drawbacks of prior synthetic methods.

Summary of the Invention

The present invention answers such a need by providing a short and regio- and stereoselective method of preparing aminosterol compounds. According to a
10 method of the invention, regio- and stereoselective oxidation and sulfonation can be achieved with fewer protecting groups and consequently fewer steps.

The invention also provides a method of regioselectively and stereoselectively oxidizing a primary hydroxyl substituent in the presence of a secondary hydroxyl substituent attached to the same fused ring system.

15 The invention further provides a method of regioselectively sulfonating one secondary hydroxyl substituent over another secondary hydroxyl substituent attached to the same fused ring system.

A method of the invention also provides novel intermediate compounds.

Brief Description Of The Drawings

20 Advantageous aspects of the invention will be evident from the following detailed description which should be considered in conjunction with the attached drawings, wherein:

Fig. 1 illustrates the chemical structure of squalamine; and


25 Fig. 2 illustrates the chemical structure of compound 1436.

Detailed Description of the Invention

Microbial hydroxylation has been achieved in steroid chemistry. Mahato, S.B., *et al.*, *Steroids*, 62, 332-345 (1997). Despreaux has described the microbial
30 7 α -hydroxylation of 3-ketobisnorcholesterol (1, Scheme 1 below) using the species *Botryodiplodia theobromae*. Despreaux, C.W., *et al.*, *Appl. Environ. Microbiol.*, 51,

946-949 (1986); Despreaux *et al.*, U.S. Patent No. 4,230,625; and Despreaux *et al.*, U.S. Patent No. 4,301,246. This invention uses steroid compound 2 as a starting material for the synthesis of squalamine, 1436 and homologous aminosterols. A method of the invention introduces the 7- α -hydroxyl group using microbial hydroxylation and proceeds without protection of the 7-hydroxyl group. A general outline of a method of the invention is outlined in Scheme 1 below:


Scheme 1



According to a method of the invention, steroid 2 may be converted to aminosterol compounds such as, but not limited to, squalamine, compound 1436 and aminosterol homologs by means of two regioselective reactions without the use of protecting groups. According to the invention, in a fused ring system, a primary hydroxyl moiety can be selectively oxidized over a secondary hydroxyl moiety. For example, if the fused ring system has a steroidal structure, as described below, a

C-22 primary hydroxyl moiety can be selectively oxidized over a secondary hydroxyl moiety at the C-7 position. Also according to the invention, in a fused ring system, one secondary hydroxyl moiety can be selectively sulfonated over another secondary hydroxyl moiety. For example, if the fused ring system has a steroidal structure, as described below, a C-24 secondary hydroxyl moiety can be selectively sulfonated over a C-7 secondary hydroxyl moiety. According to the invention, relatively high yields (e.g. 77%) as well as regioselectivity and stereoselectivity may be achieved. Some C24 selectivity has been shown in the sulfonation reaction on a spermidinyl-steroidal diol. However, this reaction not only required heating and protection of the C7-OH group, but the yield of the compound was low (10%). Moriarty, R.M., *et al.*, *Tetrahedron Lett.*, 35, 8103-8106 (1994).

An example of the invention provides a short and regioselective method of preparing an aminosterol compound of the general formula I:



In formula I, NR_1R_2 may be any saturated or unsaturated, linear or branched amino group. According to the invention, such an amino group may contain more than one nitrogen. Preferably, in formula I:

R_1 and R_2 are independently selected from the group consisting of: H, alkyl, alkenyl, alkynyl, $-(\text{CH}_2)_n\text{-NH-}(\text{CH}_2)_m\text{-NH}_2$, and $-(\text{CH}_2)_n\text{-NH-}(\text{CH}_2)_m\text{-NH-}(\text{CH}_2)_p\text{-NH}_2$;

n is an integer from 1-3;

m is an integer from 1-4; and

p is an integer from 1-2.


7

Most preferably, the compound of formula (I) is squalamine or compound 1436.

According to the invention, an aminosterol compound of formula I may be prepared by


(a) reacting compound 2:

5



2


under conditions sufficient to form compound 3:



3

10


(b) reacting compound 3 under conditions sufficient to form compound 4:



4

8

(c) reacting compound 4 under conditions sufficient to form compound 5:




5

;

(d) reacting compound 5 under conditions sufficient to form compound 7:


5



7

;

(e) reacting compound 7 under conditions sufficient to form compound 8:




8

;

(f) reacting compound 8 under conditions sufficient to form compound 9:


9



9

;


(g) reacting compound 9 under conditions sufficient to form compound 10:



10

;

(h) reacting compound 10 under conditions sufficient to form compound 11:



11

; and

5

(i) reacting compound 11 under conditions sufficient to form an aminosterol compound of the general formula (I), as described above. Each of the compounds produced by a method of the invention may be isolated and purified using techniques known in the art including, but not limited to, extraction and

chromatography. Each of the compounds produced by a method of the invention may be characterized using techniques known in the art such as, for example, mass spectrometry, ^1H NMR and ^{13}C NMR.


As set forth above, a method according to the invention includes processes
5 for regioselectively oxidizing a C-22-OH group in the presence of a C-7-OH group as well as the regioselective sulfonation of a C-24-OH group in the presence of a C-7-OH group. With respect to steps (a)-(i) of a method of the invention, "under conditions sufficient" may be any synthetic method that achieves the desired transformation without effecting the stereochemistry of the remainder of the
10 molecule. With respect to step (a), compound 2 may be transformed or converted to compound 3 using reduction methods known in the art. Despreaux, C.W., *et al.*, *Appl. Environ. Microbiol.*, 51, 946-949 (1986); Starr, J.E., Editor: C. Djerassi, Holden-Day, Inc., San Francisco, Chapter 7, pgs. 300-307 "Steroid Reaction" (1963). Preferably, reduction is achieved using lithium in ammonia with, preferably,
15 yields of at least about 76%.

Compound 3 may be transformed or converted to compound 4 by any protecting method known in the art, preferably, by ketalization of the carbonyl moiety. Ketalization may be performed utilizing ethylene glycol in chlorotrimethylsilane in good yield. Chan, T.H., *et al.*, *Synthesis*, 203-205 (1983).

20 Compound 4 may be transformed or converted to compound 5 by regioselective oxidation of the primary alcohol at the C-22 position, preferably by reaction with bleach in the presence of a catalyst. The bleach may be any bleach, preferably sodium hypochlorite (NaOCl). The catalyst may be any catalyst which in combination with the bleach achieves the regioselective oxidation. Preferably, the
25 catalyst is a TEMPO catalyst (2,2,6,6-tetramethyl-1-piperidinyloxy free radical, commercially available from Aldrich Chemicals, Milwaukee, WI). Preferably, conditions are chosen such that yields of about 98% are achieved. Anelli, P.L., *et al.*, *Org. Syn.*, Vol. 69, page 212, "A General Synthetic Method for the Oxidation of Primary Alcohols to Aldehydes: (S)-(+)-2 Methylbutanal".

30 Compound 5 may be transformed or converted to compound 7 by a carbon-carbon double bond formation reaction (e.g., Wittig reaction, Wadsworth-Emmons

reaction, Peterson olefination reaction). Preferably, compound 5 is reacted with Wadsworth-Emmons reagent 6 (Jones, S.R., *et al.*, *J. Org. Chem.*, 63, 3786-3789 (1998)):



5 to afford enone compound 7 efficiently (82%).


Compound 7 may be transformed or converted to compound 8 by reduction of the C-24 carbonyl moiety in good yield. Compound 8 may be transformed or converted to compound 9 by reduction of the C22 double bond. Preferably, reduction was achieved by means of hydrogenation. Compound 9 may be
10 transformed or converted to compound 10 by deprotection of the C3 carbonyl.

Compound 10 may be transformed or converted to compound 11 by regioselective sulfonation of C24 hydroxyl group, preferably, by reacting compound 10 with a very small excess (5%) of sulfur-trioxide complex. Preferably, the diastereomeric excess in the sulfate is about 95% based on the HPLC method.


15 Lastly, compound 11 may be transformed or converted to the desired aminosterol compound (e.g. squalamine, compound 1436 or homologous compounds) by any means whereby a carbonyl moiety may be converted to an amino group including, but not limited to, reductive amination conditions. Rao, M., *et al.*, *J. Nat. Prod.* 63, pp. 631-635 (2000); Zhang, X., *et al.*, *J. Org. Chem.* 63,
20 8599-8603 (1998); and Weis, A.L., *et al.*, *Tetrahedron Lett.*, 40, 4863-4864 (1999).

An example of a preferred method of preparing aminosterol compound squalamine is illustrated in Scheme 2 below:


Scheme 2



The invention also provides a method of regioselectively oxidizing a primary hydroxyl substituent in the presence of a secondary hydroxyl substituent attached to the same fused ring base. According to this embodiment of the invention, a fused ring base to which both a primary hydroxyl substituent and a secondary hydroxyl substituent are attached is reacted with bleach in the presence of a catalyst whereby solely the primary hydroxyl substituent is oxidized to an aldehyde. According to the invention a fused ring base is any compound containing at least two saturated and/or unsaturated ring systems which share at least two carbon atoms. According to the invention, the fused ring base may also contain appropriate substituents (e.g. alkyl groups, hydroxyl groups, amino groups, etc.) or unsaturations (e.g. double bonds, triple bonds, carbonyl groups). An appropriate substituent or unsaturation is one that would not adversely effect the desired transformation or conversion, as described below. Preferably, the fused ring base is a steroid ring system having the following general formula:




where R is a linear or branched, substituted or unsubstituted, saturated or unsaturated alkyl group. Preferably, the fused ring base has one of the following structures:




and

14




The bleach and the catalyst are each as described herein.

The invention also provides for a method of regioselectively sulfonating one secondary hydroxyl substituent in the presence of another secondary hydroxyl substituent attached to the same fused ring base. The fused ring base is as described above except that the preferred fused ring base has the following structure:



According to this embodiment of the invention, a fused ring base to which two secondary hydroxyl substituents are attached is reacted with sulfur-trioxide pyridine complex (commercially available from Aldrich Chemical, Milwaukee, WI):



The methods of the invention achieve regioselectivity of one hydroxyl moiety in the presence of another unprotected hydroxyl moiety. The methods of the invention achieve regioselectivity of at least about 9:1 excess of the desired hydroxylated or sulfonated compound. Preferably, selectivity of greater than about 19:1 is achieved, and most preferably, greater than about 33:1 selectivity is

achieved.

The methods of the invention as described above may be used to produce a hydroxylated intermediate that can be further modified, as described above, to produce the desired final product. The methods of the invention produce
5 regiospecific intermediates that can be further modified to synthesize squalamine, compound 1436, other useful aminosterols or steroids having stereospecific groups (e.g., C-24 sulfate groups in an *R* orientation for, squalamine and compound 1436). Such intermediates include, but are not limited to, compounds 3-10 as illustrated in Scheme 2 above.

10 The methods of the invention will now be described in specific examples. However, the following examples serve merely to illustrate the invention and are not meant to limit the invention in any manner.

Examples

15 Regioselective and Stereoselective Synthesis of a Precursor for Squalamine, Compound 1436 or Homologous Aminosterols

General. The ^1H and ^{13}C NMR spectra were generated at 400 and 100 MHz, utilizing 7.28 and 77.0 (CDCl_3) ppm as the references respectively. Elemental
20 analyses were performed at Oneida Research Services, Inc., Whitesboro, NY. Fast Atom Bombardment mass spectral analysis was carried out at M-Scan Inc., West Chester, PA.

Example 1. Preparation of (5- α -, 7- α)-3-Ketobisnorcholan-7,22-diol (3).

25 Liquid ammonia (125 mL) was treated with tetrahydrofuran (15 mL) and lithium (300 mg, 43 mmol) and stirred for 30 min. Then a solution of 2 (Despreaux, C.W., *et al.*, *Appl. Environ. Microbiol.*, 51, 946-949 (1986)) (352 mg, 1.20 mmol) in tetrahydrofuran (20 mL) and ethanol (0.4 mL) was added. The reaction mixture was stirred for 40 min and then 20 g of ammonium chloride was
30 added. The solvent was evaporated under nitrogen and the residue was treated with water (200 mL) and extracted with ethyl acetate (3 x 75 mL). The organic phase was

washed with brine, dried over sodium sulfate, filtered, and evaporated. Purification of the resulting solid by flash chromatography on silica gel (hexane-ethyl acetate-methanol 10:10:1) afforded pure **3** (251 mg, 71%, mp 221-223 °C, MW 348.53); ¹H NMR (CDCl₃): δ 3.86 (br s, 1H), 3.65-3.62 (m, 1H), 3.39-3.36 (m, 1H), 2.34-1.18 (m, 23H), 1.05 (d, J = 6.6 Hz, 3H), 1.01 (s, 3H), 0.71 (s, 3H); ¹³C NMR (CDCl₃): δ 67.9, 67.4, 52.4, 50.2, 45.2, 44.1, 42.7, 39.5, 39.2, 39.0, 38.7, 38.1, 36.5, 35.6, 27.7, 23.7, 21.2, 16.7, 11.9, 10.4; MS (+FAB): 349 ([M+I]⁺, 100), 331 (52); Anal. Calcd for C₂₂H₃₆O₃: C, 75.82; H, 10.41. Found: C, 75.71; H, 10.19.

Example 2. (5- α -, 7- α)-3-Dioxolane Bisnorcholan-7,22-diol (4).

To a mixture of steroid **3** (101g, 0.290 mol) of Example 1 and anhydrous ethylene glycol (800 mL) was added chlorotrimethylsilane (200 mL, 1.58 mol) over 60 min at room temperature under nitrogen. The reaction mixture was stirred at room temperature for 19 h. The mixture was poured slowly into saturated sodium bicarbonate solution (1 L) and extracted with dichloromethane (3 x 500 mL). The organic layer was washed with brine (3 x 150 mL) and dried over sodium sulfate (20 g). After filtration and evaporation, the product was recrystallized from ethyl acetate in hexane (800 mL). The solid was filtered and washed with hexane (150 mL) to afford **4** (96.14 g, 84%, mp 173-175 °C, MW 392.58); ¹H NMR (CDCl₃): δ 3.93 (s, 4H), 3.83 (br s, 1H), 3.65 (d of d, J = 10.4 and 3.1 Hz, 1H), 3.36 (d of d, J = 10.4 and 7.1 Hz, 1H), 2.0-1.8 (m, 3H), 2.7-1.1 (m, 21H), 1.05 (d, J = 6.6 Hz, 3H), 0.82 (s, 3H), 0.69 (s, 3H); ¹³C NMR (CDCl₃): δ 109.2, 67.8, 64.1, 52.4, 50.3, 45.6, 42.7, 39.5, 39.3, 38.8, 37.4, 36.2, 36.1, 35.7, 35.5, 31.2, 27.7, 23.7, 20.9, 16.7, 11.9, 10.3; MS (+FAB): 394 ([M+I]⁺, 100); Anal. Calcd for C²⁴H⁴⁰O₄: C, 73.43; H, 10.27. Found: C, 73.15; H, 10.15. This reaction was accomplished at 10% concentration of substrate, which allows for efficient scale-up of the procedure.

Example 3. Preparation of (5- α -, 7- α)-3-Dioxolane-7-hydroxy Bisnorcholan-22-al (5).

To a solution of **4** (100 g, 255 mmol) of Example 2 in methylene chloride (1,200 mL) was added potassium bromide (3.19g, 26.8 mmol) and sodium

bicarbonate (10.97g, 130 mmol) dissolved in water (120 mL). The cooled (0 °C) reaction mixture was treated with TEMPO (1.20 g, 7.7 mmol) and 10-13% sodium hypochlorite (170 mL, 275-358 mmol). After stirring (magnetic) for 2 h at 0 °C, the reaction mixture was treated with sodium thiosulfate (20 g, 126 mmol) in water (220 mL). The organic phase was separated, washed with brine (3 x 70 mL), dried over sodium sulfate (30 g), filtered, and concentrated *in vacuo* for 18 h at room temperature to afford **5** (99.5 g, 98%, MW 390.57, FW 397.77); ¹H NMR (CDCl₃): δ 9.57 (d, J = 3.4 Hz, 1H), 3.95 (s, 4H), 3.83 (br s, 1H), 3.76 (m, 1H), 2.35 (m, 1H), 2.0-1.2 (m, 21 H), 1.13 (d, J = 6.8 Hz, 3H), 0.83 (s, 3H), 0.72 (s, 3H); ¹³C NMR (CDCl₃): δ 204.9, 109.0, 67.6, 64.0, 50.8, 49.7, 49.3, 45.4, 43.0, 39.3, 39.0, 37.3, 36.2, 35.9, 35.6, 35.4, 31.0, 26.8, 23.8, 20.7, 13.3, 12.1, 10.2; MS (+FAB): 391 ([M+H]⁺, 100); Anal. Calcd for C₂₄H₃₈O₄·H₂O: C, 72.47; H, 9.83. Found: C, 72.49; H, 9.77.

Example 4. Preparation of (5- α -, 7- α)-3-Dioxolane-7-hydroxy Cholest-23-en-24-one (**7**).

A mixture of 97% sodium *t*-butoxide (37 g, 373 mmol) and anhydrous tetrahydrofuran (400 mL) was stirred for 10 min under nitrogen and then a solution of **6** (94 g, 423 mmol, see Scheme 2 above) in tetrahydrofuran (150 mL) was added in one portion. The mixture initially warmed to 41 °C, but returned to 24 °C while stirring (45 min). Then a solution of **5** (99.48 g, 250 mmol) of Example 3 in tetrahydrofuran (400 mL) was added over 60 min. The reaction mixture was stirred overnight at room temperature (18 h) and then water was added (30 mL). The reaction mixture was concentrated *in vacuo* and treated with cyclohexane (1200 mL), toluene (600 mL) and water (160 mL). The organic layer was separated, washed with brine (3 x 100 mL) and water (160 mL), dried over sodium sulfate (30 g), filtered, and evaporated to yield a solid. The crude solid was recrystallized from ethyl acetate in cyclohexane and dried *in vacuo* at 50 °C for 5 h to yield **7** (94.64 g, 82%, mp 177-178 °C, MW 458.69); ¹H NMR (CDCl₃): δ 6.72 (d of d, J = 15.7 and 9.0 Hz, 1H), 6.07 (d, J = 15.7 Hz, 1H), 3.94 (s, 4H), 3.83 (br s, 1H), 2.85 (hept, J = 6.9 Hz, 1H), 2.29 (m, 1H), 2.0-1.1 (m, 22 H), 1.11 (m, 9H), 0.83 (s, 3H), 0.71 (s,

3H); ^{13}C NMR (CDCl_3): δ 204.5, 152.4, 126.2, 109.1, 67.8, 64.1, 54.9, 50.4, 45.6, 43.0, 40.0, 39.5, 39.3, 38.1, 37.4, 36.3, 36.1, 35.7, 31.2, 28.1, 23.6, 20.9, 19.3, 18.6, 18.4, 12.1, 10.3; MS (+FAB): 459 ($[\text{M}+1]^+$, 92), 99 (100); Anal. Calcd for $\text{C}_{29}\text{H}_{46}\text{O}_4$: C, 75.94; H, 10.11. Found: C, 75.57; H, 9.87.

5

Example 5. Preparation of (5- α -, 7- α -, 24S-)-7, 24-Dihydroxy-3-dioxolane Cholest-23-ene (8).

A dried and nitrogen blanketed reactor was charged with 1 M (R)-MeCBS reagent in toluene (20 mL, 20 mmol) and 1 M borane-tetrahydrofuran complex in
10 tetrahydrofuran (25 mL, 25 mmol) and stirred for 2 h at room temperature. The reaction mixture was cooled (-15 to -28°C), treated with steroid 7 (9.16 g, 20 mmol) of Example 4 in tetrahydrofuran (150 mL), and stirred for 2 hr (-20 to -28 °C). The reaction mixture was treated with methanol (25 mL) with stirring for 18 hr at room temperature, and then repeatedly evaporated by distillation and treated with
15 methanol (4 x 30 mL) to exchange solvents. Finally methanol (70 mL) was added and the reaction mixture was brought to reflux, cooled in the freezer (no crystals formed), and concentrated *in vacuo*. Recrystallization from acetonitrile (100 mL), filtration, and evaporation at 50-60 °C for 7 hr afforded crystals of 8 (7.43 g, 80%, mp 121-125 °C, MW 460.70, FW 464.31); ^1H NMR (CDCl_3): δ 5.5-5.3 (m, 2H),
20 3.94 (s, 4H), 3.82 (br s, 1H), 3.75 (in, 1H), 2.2-1.1 (m, 25H), 1.05 (d, J = 6.6 Hz, 3H), 0.94 (d, J = 6.7 Hz, 3H), 0.88 (d, J = 6.8 Hz, 3H), 0.83 (s, 3H), 0.70 (s, 3H); ^{13}C NMR (CDCl_3): δ 139.5, 128.6, 109.2, 78.5, 67.8, 64.1, 55.5, 50.6, 45.6, 42.6, 40.0, 39.5, 39.4, 37.5, 36.2, 36.1, 35.7, 35.6, 33.9, 31.2, 28.7, 23.6, 20.9, 20.4, 18.3, 18.1, 12.0, 10.3; MS (+FAB): 462 ($[\text{M}+1]^+$, 100); Anal. Calcd for $\text{C}_{29}\text{H}_{48}\text{O}_4 \cdot 0.2\text{H}_2\text{O}$: C, 75.02; H, 10.51. Found: C, 75.00; H, 10.48.
25

Example 6. Preparation of (5- α -, 7- α -, 24R-)-7, 24-Dihydroxy-3-dioxolane Cholestane (9).

Steroid 8 (10.0 g, 21.5 mmol) of Example 5, toluene (170 mL), triethylamine
30 (1 mL), and 10% platinum on carbon (0.5 g) were combined under 50 psi of hydrogen in a Parr apparatus (19 h). The reaction mixture was filtered through Celite

(10 g), washed with chloroform and ethyl acetate (10 mL total), and concentrated *in vacuo* to afford a solid, which was recrystallized from ethyl acetate in hexane (180 mL). The solid was filtered and concentrated at 50-60 °C under vacuum for 7 h to afford pure **9** (9.24 g, 92%, mp 161-163 °C, MW 462.72, FW 466.32); ¹H NMR (CDCl₃): δ 3.95 (s, 4H), 3.84 (br s, 1H), 3.33 (br s, 1H), 2.0-1.1 (m, 29H), 0.93 (m, 9H), 0.83 (s, 3H), 0.67 (s, 3H); ¹³C NMR (CDCl₃): δ 109.2, 77.0, 67.8, 64.1, 55.9, 50.5, 45.5, 42.6, 39.5, 37.4, 36.2, 36.1, 35.7, 35.5, 33.5, 32.0, 31.2, 30.5, 28.2, 23.6, 20.9, 18.8, 18.6, 17.2, 11.8, 10.3; MS (+FAB): 463 ([M+I]⁺, 100)⁺, Anal. Calcd for C₂₉H₅₀O₄-0.2H₂O: C, 74.70; H, 10.89. Found: C, 74.48; H, 10.49.

Example 7. Preparation of (5- α -, 7- α -, 24*R*)-7, 24-Dihydroxy-3-ketocholestane (**10**).

Steroid **9** (2.03 g, 4.35 mmol) of Example 6, *p*-toluenesulfonic acid (200 mg), water (1 mL), and acetone (100 mL) were combined with stirring for 4 h. The reaction mixture was concentrated *in vacuo* and treated with dichloromethane (100 mL) and saturated sodium bicarbonate solution (50 mL). The organic layer was removed, washed with brine (3 x 25 mL), dried over sodium sulfate (10 g), filtered, and evaporated at 50-60 °C. The solid was recrystallized from ethyl acetate in hexane (50 mL), filtered, washed with hexane, and dried *in vacuo* at 50-60 °C for 7 hr to afford **10** (1.63 g, 89%, mp 151-153 °C, MW 418.67); ¹H NMR (CDCl₃): δ 3.88 (br s, 1H), 3.33 (br s, 1H), 2.5-1.1 (m, 29H), 1.02 (s, 3H), 0.94 (m, 9H), 0.71 (s, 3H); ¹³C NMR (CDCl₃): δ 212.0, 76.9, 67.3, 56.1, 50.3, 45.1, 44.1, 42.6, 39.4, 39.0, 38.1, 38.0, 36.6, 35.8, 35.6, 33.6, 32.1, 30.6, 28.2, 23.6, 21.1, 18.9, 18.6, 17.3, 11.8, 10.4; MS (+FAB): 419 ([M+I]⁺, 100); Anal. Calcd for C₂₇H₄₆O₃: C, 77.46; H, 11.07. Found: C, 77.25; H, 11.04.

Example 8. Preparation of Potassium salt of (5- α -, 7- α -, 24*R*)-7-Hydroxy-3-ketocholestan-24-yl Sulfate (**11**).

A dried and nitrogen blanketed flask was treated with compound **10** (2.09 g, 5.0 mmol) of Example 7 dissolved in anhydrous pyridine (30 mL). Sulfur trioxide pyridine complex (836 mg, 5.25 mmol, 1.05 equiv.) dissolved in pyridine (20 mL)

was added to the reaction mixture, which was stirred for 4 h at room temperature. Water was added (10 mL) and the pyridine was removed by concentration *in vacuo* at 40 °C. The residue was treated with ethyl acetate (50 mL) and potassium chloride (1.12 g, 15 mmol) dissolved in water with stirring for 1.5 h. The potassium salt of 11
5 was collected on Celite (3 g) by filtration, washed with ethyl acetate (50 mL) and water (10 mL), and dissolved in 1 N potassium hydroxide in 15 methanol (10 mL, 10 mmol) and methanol (100 mL). The methanol was removed *in vacuo* to dryness and the solid was washed with water (30 mL), filtered, and dried *in vacuo* at room temperature for 20 hr to afford 11 (2.10 g, 77%, MW 536.82, FW 544); ¹H and ¹³C
10 NMR were identical to published spectra. HPLC analysis by the method described previously (Zhang, X., *et al.*, *J. Org. Chem.*, 63, 8599-8603 (1998)) indicated a diastereomeric excess of 95 %.

Example 9. Preparation of Compound 1436

15 A clear colorless solution of compound 11 (16 mg, 0.032 mmol) and spermine (20 mg, 0.1 mmol, commercially available from Aldrich) in anhydrous methanol (3 ml) was stirred at room temperature under nitrogen for 12 hours, cooled to -78°C, and treated dropwise with sodium borohydride (1 pellet, 0.4 g, 10 mmol) in methanol (10 ml). This reaction mixture was stirred for 3 hours, treated with a
20 mixture of water and methanol (10 ml each), warmed to room temperature, and then treated with 0.78% trifluoroacetic acid (TFA) solution until its pH reached the range of 4 - 5. The resulting mixture was filtered through a thin pad of Celite®, and the Celite® was washed with methanol and water (100 ml). Celite® is SiO₂ that is commercially available from Aldrich. The combined acidic washes were
25 concentrated *in vacuo* at room temperature and then freeze-dried overnight to give a white solid. The Celite® cake was then washed with isopropyl amine/methanol/water (140 ml of 1:3:3), and the basic portion was evaporated to reduce its volume. This material was freeze-dried overnight to give a light brown solid. Both washes contained compound 1436, so they were combined and acidified
30 to a pH of 3 with 0.78% TFA, filtered, and loaded onto a small HPLC column (1 cm diameter, see below). The reaction product was compound 1436 (12.2 mg, 36%):

¹H NMR (400 MHz, D₂O): δ 4.14 (m, 1H), 3.83 (m, 1H), 3.2 - 3.0 (m, 13H), 2.1-1.0 (m, 35H), 0.92 (m, 9H), 0.82 (s, 3H), 0.67 (s, 3H); ¹³C NMR (400 MHz, D₂O): δ 87.2, 68.0, 57.9, 56.0, 50.5, 47.4, 45.6, 44.9, 42.8, 41.9, 39.7, 37.5, 36.9, 36.7, 36.0, 35.8, 31.5, 31.1, 30.6, 28.3, 27.1, 24.8, 24.1, 23.6, 23.4, 23.1, 21.4, 19.2, 17.7, 12.1, 11.2; MS (-LD): 684 (M-1); Anal. Calcd. for C₃₇H₇₂N₄O₅S-3TFA-2H₂O: C, 48.58; H, 7.49; F, 16.08; N, 5.27; S, 3.02. Found: C, 48.49; H, 7.40; F, 16.16; N, 5.31; S, 3.05.

Example 10. Purification of Compound 1436 by HPLC

The crude material of Example 9 was dissolved in water (50 ml), cooled in an ice bath, and acidified with 1.5% TFA in water until its pH was 3. Initially, it was observed that one obtains a suspension as the pH drops, and then a solution is obtained at lower pH. This solution was loaded onto a Rainin reverse phase HPLC system (2.14 cm diameter, C18, 100 Å, 8µm) and eluted with A (water with 0.1% TFA) and B (acetonitrile with 0.1% TFA). The HPLC program was as follows: 10 min (0 - 10% B), 60 min (10-45% B), 10 min (45-80% B), 10 min (80% B). Pure product eluted in the 33 to 55 minute fractions, as determined by TLC (R_f: 0.1-0.2 in 6/3/1 CH₂Cl₂/MeOH/NH₄OH)(should evaporate plates under vacuum before eluting, and observe with ninhydrin stain after eluting), which was lyophilized to produce 1.20 grams of compound 1436 as a white powder (70%); C₃₇H₇₂N₄O₅S-3TFA-2.5H₂O, FW 1072.18).

Example 11. Preparation of Squalamine

Squalamine was prepared by reacting the potassium salt of compound 11 (0.5 equivalents) of Example 8 with H₂N(CH₂)₃NH(CH₂)₄N₃ • 2HCl (1 equivalent) in NaOMe (2 equiv) and methanol at room temperature for 24 hours and then at -78°C with NaBH₄ followed by treatment with H₂, RaNi, RP-HPLC, 69% based on the potassium salt of compound 11. See Weis *et al.*, Tetrahedron Letters, 40, 4863-4864 (1999).

In describing the invention, applicant has stated certain theories in an effort to disclose how and why the invention works in the manner in which it works. These theories are set forth for informational purposes only. Applicants do not wish to be bound by any specific theory of operation.

5

While the invention has been described in terms of various specific preferred embodiments and specific examples, those skilled in the art will recognize that various changes and modifications can be made without departing from the spirit and scope of the invention, as defined in the appended claims.

10

Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the illustrative examples, make and utilize the compounds of the present invention and practice the claimed methods. It should be understood that the foregoing discussion and examples merely present a detailed description of certain preferred embodiments. It will be apparent to those of

15

ordinary skill in the art that various modifications and equivalents can be made without departing from the spirit and scope of the invention. All the patents, journal articles and other documents discussed or cited above are herein incorporated by reference in their entirety.


20

WHAT IS CLAIMED:

1. A method for regioselectively oxidizing a primary hydroxyl substituent attached to a fused ring base in the presence of a secondary hydroxyl substituent attached to said fused ring base comprising the step of:

5 reacting a fused ring system comprising a primary hydroxyl substituent and a secondary hydroxyl substituent with bleach in the presence of a catalyst wherein solely said primary hydroxyl substituent is oxidized to an aldehyde.

2. A method of claim 1, wherein said fused ring system is a steroid ring structure of the general formula:



where R is a linear or branched, substituted or unsubstituted, saturated or unsaturated alkyl group.


15

3. A method of claim 2, wherein the base structure of said fused ring system is substituted with at least one hydroxyl group to form a secondary hydroxyl group and where R is a linear or branched, saturated or unsaturated alkyl chain substituted with at least one hydroxyl group to form a primary hydroxyl group.

20


4. A method of claim 3, wherein said fused ring structure has the following formula:

24



5. A method of claim 4, wherein said fused ring system has the following structure:


5



6. A method for regioselectively sulfonating a first secondary hydroxyl
 10 substituent attached to a fused ring base in the presence of a second secondary
 hydroxyl substituent attached to said fused ring base comprising the step of:
 reacting a fused ring system comprising a first secondary hydroxyl
 substituent and a second secondary hydroxyl substituent with a sulfur trioxide-
 pyridine complex to regioselectively sulfonate said first secondary hydroxy
 15 substituent.

7. A method of claim 6, wherein said fused ring system is a steroid ring
 structure of the general formula:

25




where R is a linear or branched, substituted or unsubstituted, saturated or unsaturated alkyl group.

5

8. A method of claim 7, wherein R is a linear or branched, saturated or unsaturated alkyl chain substituted with at least one hydroxyl group to form said first secondary hydroxyl group and where said fused ring system is substituted with at least one hydroxyl group to form said second secondary hydroxyl group.

10


9. A method of claim 8, wherein said fused ring structure has the following formula:



15 10. A method for preparing an aminosterol compound comprising the steps of:


(a) reacting compound 2:

26



2

under conditions sufficient to form compound 3:




3

5

;

(b) reacting compound 3 under conditions sufficient to form compound 4:




4

;

10

(c) reacting compound 4 under conditions sufficient to form compound 5:


27



5

;

(d) reacting compound 5 under conditions sufficient to form compound 7:




7

;

5

(e) reacting compound 7 under conditions sufficient to form compound 8:




8

;

(f) reacting compound 8 under conditions sufficient to form compound 9:


28



9

;


(g) reacting compound 9 under conditions sufficient to form compound 10:



10

;

(h) reacting compound 10 under conditions sufficient to form compound 11:




11

; and

(i) reacting compound 11 under conditions sufficient to form an aminosterol compound of the general formula:

29



where NR_1R_2 forms a saturated or unsaturated, linear or branched amino group.

- 5 11. A method of claim 10, wherein R_1 and R_2 are independently selected from the group consisting of: H, alkyl, alkenyl, $-(CH_2)_n-NH-(CH_2)_m-NH_2$, and $-(CH_2)_n-NH-(CH_2)_m-NH-(CH_2)_p-NH_2$;
 n is an integer from 1-3;
 m is an integer from 1-4; and
10 p is an integer from 1-2.
12. A method of claim 10, wherein said conditions sufficient comprise reacting said compound 4 with bleach in the presence of a catalyst to form said compound 5.
- 15 13. A method of claim 12, wherein said conditions sufficient comprise reacting said compound 10 with a sulfur-trioxide pyridine complex.
14. A method of claim 13, wherein said aminosterol compound is squalamine or compound 1436.
- 20 15. A compound selected from the group consisting of:




FIG. 1





FIG. 2



INTERNATIONAL SEARCH REPORT

Inter il Application No
PCT/US 01/12004

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07J9/00 C07J21/00 C07J41/00 C07J31/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07J

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, EPO-Internal, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DATABASE CHEMABS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; DESPREAUX, CARL W. ET AL: "Microbial 7.alpha.-hydroxylation of 3-ketobisnorcholesterol" retrieved from STN Database accession no. 105:21428 XP002175382 abstract & APPL. ENVIRON. MICROBIOL. (1986), 51(5), 946-9 ,	15
X	EP 0 018 515 A (HOFFMANN LA ROCHE) 12 November 1980 (1980-11-12) example 1 ----- -/-	15

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

Z document member of the same patent family

Date of the actual completion of the international search

21 August 2001

Date of mailing of the international search report

10/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Watchorn, P

INTERNATIONAL SEARCH REPORT

Inter if Application No
PCT/US 01/12004

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 98 24800 A (MAGAININ PHARMA) 11 June 1998 (1998-06-11) figures 15A, 15B; examples 24, 24B, 26-28, 28B, 29, 30, 30B, 31, 31B, 32	1-15
Y	XUEHAI ZHANG ET AL: "Synthesis of Squalamine Utilizing a Readily Accessible Spermidine Equivalent" JOURNAL OF ORGANIC CHEMISTRY., vol. 63, no. 23, 1998, pages 8599-8603, XP002175377 AMERICAN CHEMICAL SOCIETY. EASTON., US ISSN: 0022-3263 page 8599, column 1, paragraph 1 page 8600, schemes 1 and 2	1-15
Y	S. R. JONES ET AL: "Efficient Route to 7.alpha.-(Benzoyloxy)-3-dioxolane Cholestan-24(R)-ol, a Key Intermediate in the Synthesis of Squalamine" JOURNAL OF ORGANIC CHEMISTRY., vol. 63, no. 11, 1998, pages 3786-3789, XP002175378 AMERICAN CHEMICAL SOCIETY. EASTON., US ISSN: 0022-3263 page 3788, scheme 1	1-15
Y	RAO M N ET AL: "PRACTICAL APPROACHES TO REMOTE ASYMMETRIC INDUCTION IN STEROIDAL SIDE-CHAINS UTILIZING OXAZABOROLIDINE REAGENTS" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, US, vol. 62, no. 13, 27 June 1997 (1997-06-27), pages 4541-4545, XP002062007 ISSN: 0022-3263 page 4542, scheme 2 page 4543, scheme 3	1-15
Y	WEIS A L ET AL: "Synthesis of an Azido Spermidine Equivalent" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 40, no. 26, 25 June 1999 (1999-06-25), pages 4863-4864, XP004168666 ISSN: 0040-4039 page 4864, paragraph 1	1-15

-/--

INTERNATIONAL SEARCH REPORT

Inter if Application No
PCT/US 01/12004

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	MORIARTY R M ET AL: "SYNTHESIS OF SQUALAMINE A STEROIDAL ANTIBIOTIC FROM THE SHARK" TETRAHEDRON LETTERS, ELSEVIER SCIENCE PUBLISHERS, AMSTERDAM, NL, vol. 35, no. 44, 31 October 1994 (1994-10-31), pages 8103-8106, XP000572370 ISSN: 0040-4039 page 8104, reaction of compound 16 to compound 1 page 8105, paragraph 2	1-15
Y	PECHULIS A D ET AL: "SYNTHESIS OF 23XL-SQUALAMINE, AN ANTI-INFECTIVE STEROIDAL POLYAMINE" JOURNAL OF ORGANIC CHEMISTRY, AMERICAN CHEMICAL SOCIETY. EASTON, US, vol. 60, no. 16, 11 August 1995 (1995-08-11), pages 5121-5126, XP000572371 ISSN: 0022-3263 page 5123, column 1, paragraph 2	1-15
Y	A. M. MAIONE ET AL: "Carbonyl Compounds by Osmium Tetroxide Oxidation: Preferential Oxidation of Primary over Secondary Hydroxy Groups" SYNTHESIS., no. 11, November 1984 (1984-11), pages 955-957, XP002175379 GEORG THIEME VERLAG. STUTTGART., DE ISSN: 0039-7881 page 956; table 3	1-15
P,X	KINNEY, WILLIAM A. ET AL: "A Short Formal Synthesis of Squalamine from a Microbial Metabolite" ORG. LETT. 2(19), 2921 -2922 , 21 September 2000 (2000-09-21), XP002175380 page 2922, scheme 1 -& "supporting information" ORG. LETT., 'Online! vol. 2, no. 19, 21 September 2000 (2000-09-21), XP002175381 Retrieved from the Internet: <URL:http://pubs.acs.org/subscribe/journals/orlef7/supinfo/o10059495/o10059495_s1.pdf> 'retrieved on 2001-08-21!	1-15

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 01 /2004

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 1-9 (in part)

Present process claims 1-9 relate to processes defined by reference to a desirable characteristic or property, namely in the case of claim 1, that the oxidation of the primary hydroxy group is performed using bleach with a catalyst defined in terms of it selectively facilitating the oxidation of the primary alcoholic group (no other technical details relating to the catalyst are given). In the case of claim 6 the reaction is defined in terms of a result to be achieved in that the claim specifies a regiospecific sulphation reaction whereby the reaction conditions are defined in terms of the use of a well known sulphation agent (pyridine:SO₃) used in such a way as to give a selective sulphation reaction with no technical details as to how this regioselectivity is to be achieved.

Claim 1 and claims 2-5 dependent thereon as well as claim 6 and claims 7-9 dependent thereon cover all processes having the above mentioned characteristics or properties, whereas the application provides support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for only a very limited number of such processes. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the processes by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible.

Consequently, the search has been carried out for those parts of the claims which appear to be clear, supported and disclosed, namely those parts relating to the process of claims 1-9 insofar as they are commensurate in scope with the multistep process of claim 10 (and claims 11-14 dependent thereon).

Furthermore, claim 6 specifies a regiospecific sulphation reaction, whereby a first secondary OH group of a fused ring base containing both a first and second secondary OH group is preferentially sulphated. This claim (and also claims 7 and 8 dependent thereon) does not indicate how the two secondary OH groups differ from one another and which of them is to be sulphated. This renders claim 6 highly unclear within the meaning of Article 6 PCT, since the skilled person is not only unable to ascertain how to achieve the desirable goal of preferential sulphation because he is not given the technical details of how to achieve this (see above), but also because he does not even know from these claims what the desired property actually is. This is a further reason for the above mentioned limitation of the search on the process of claims 6-8.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an

INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 01 /2004

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.

INTERNATIONAL SEARCH REPORT

Information on patent family members

Inter-Office Application No

PCT/US 01/12004

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0018515 A	12-11-1980	US 4230625 A AT 1710 T DE 3060991 D JP 56008399 A US 4301246 A	28-10-1980 15-11-1982 02-12-1982 28-01-1981 17-11-1981
WO 9824800 A	11-06-1998	US 6262283 B AU 5591498 A EP 0942918 A JP 2001505207 T	17-07-2001 29-06-1998 22-09-1999 17-04-2001